arXiv:1209.1884v1 [quant-ph] 10 Sep 2012

Measurement-Induced Nonlocality in an n-partite quantum state.

Ali Saif M. Hassan[1,*] and Pramod S. Joag[2,†]
[1]*Department of Physics, University of Amran, Amran, Yemen*
[2]*Department of Physics, University of Pune, Pune, India-411007.*
(Dated: September 11, 2012)

We generalize the concept of measurement-induced non-locality (MiN) to n-partite quantum states. We get exact analytical expressions for MiN in an n-partite pure and n-qubit mixed state. We obtain the conditions under which MiN equals geometric quantum discord in an n-partite pure state and an n-qubit mixed state.
PACS numbers: 03.65.Ud;75.10.Pq;05.30.-d

Measurement induced non-locality (MiN) is a measure of quantum correlations as manifested in the non-local effects of local (on a single part) quantum operations[1, 2]. These local quantum operations leave invariant the reduced density operators of the part on which they act, while changing the global quantum state. MiN concerns the von-Neumann measurement on a part of a quantum system. MiN being an inherently quantum phenomenon, is expected to be useful as a tool for quantitative specification of quantum correlation. Such a quantitative specification of quantum correlations in terms of MiN was given in [2] for bipartite quantum systems. Here we generalize this measure to N-partite quantum systems. MiN is a manifestation of the quantum verses classical paradigm of quantum correlations and naturally compares with quantum discord [3–6] which is also a manifestation of such a paradigm. In fact, it is quite relevant to inquire about the conditions on quantum states under which MiN and geometric discord are equal (or, rather are different) and the different kinds of information they give about the quantum correlations in a quantum state. Here we establish such general conditions in n-partite pure and n-qubit mixed states.

Multipartite generalization of MiN : Multipartite generalization of MiN can be obtained in a manner analogous to that of geometric quantum discord [6]. For an n-partite system in a state ρ we define, for (normalized) MiN [7]

$$N_l(\rho) = \frac{d_l}{d_l - 1} \max_{\Pi^{(l)}} ||\rho - \Pi^{(l)}(\rho)||^2), l = 1, 2, \cdots, n \tag{1}$$

where $\Pi^{(l)} = \{\Pi_k^{(l)}\}$ stands for the set of von-Neumann measurements on the lth part such that $\Pi^{(l)}(\rho^{(l)}) = \sum_k \Pi_k^{(l)} \rho^{(l)} \Pi_k^{(l)} = \rho^{(l)}$, $\rho^{(l)}$ being the reduced density operator obtained by tracing out all parts other than the lth part from the n-partite state acting on $\mathcal{H} = \mathcal{H}^1 \otimes \mathcal{H}^2 \otimes \cdots \otimes \mathcal{H}^n$ with $dim(\mathcal{H}^m) = d_m,\ m = 1, 2, \cdots, n$. Such a measurement $\Pi^{(l)}$ is defined by the projectors corresponding to the eigenstates of $\rho^{(l)}$. When all the eigenvalues of $\rho^{(l)}$ are non-degenerate, there is only one von-Neumann measurement $\Pi^{(l)}$ satisfying $\Pi^{(l)}(\rho^{(l)}) = \sum_k \Pi_k^{(l)} \rho^{(l)} \Pi_k^{(l)} = \rho^{(l)}$ and the maximization requirement in Eq.(1) drops out. If one or more eigenvalues of $\rho^{(l)}$ are degenerate, the right hand side of Eq.(1) has to be maximized over the eigenspaces of degenerate eigenvalues, which is, in general, a difficult task.

Throughout this letter the superscript t denotes the transpose of a vector or a matrix.

Comparing the definitions of MiN $N_l(\rho)$ and the geometric discord $D_l(\rho)$ [6] it follows that, for any n-partite state, $N_l(\rho) \geq D_l(\rho)$. We are interested in finding the criteria for their equality.

The multipartite non-locality can be evaluated for an n-partite pure state via the following

---

[*] Electronic address: alisaif73@gmail.com
[†] Electronic address: pramod@physics.unipune.ac.in

Theorem 1: Let $|\psi\rangle = \sum_{i_1 i_2 \cdots i_n} a_{i_1 i_2 \cdots i_n} |i_1 i_2 \cdots i_n\rangle$ be a n-partite pure state. Then

$$N_l(|\psi\rangle\langle\psi|) = \frac{d_l}{d_l - 1}(1 - tr(\rho^{(l)})^2). \tag{2}$$

where $\rho^{(l)}$ is the reduced density matrix of the lth part snd $d_l = dim(H^{(l)})$.

Proof: In order to get $N_l(|\psi\rangle\langle\psi|)$ we can directly calculate the terms which define it (Eq.(1)). We have

$$\rho = |\psi\rangle\langle\psi| = \sum_{i_1 i_2 \cdots i_n} \sum_{j_1 j_2 \cdots j_n} a_{i_1 i_2 \cdots i_n} a^*_{j i_1 j_2 \cdots j_n} |i_1 i_2 \cdots i_n\rangle\langle j_1 j_2 \cdots j_n|$$

The set of von-Neumann measurements on the lth part is given by

$$\Pi^{(l)} = \{\Pi_k^{(l)} = U|k_l\rangle\langle k_l|U^\dagger\}$$

where $\{|k_l\rangle\}$, $k_l = 1, \ldots, d_l = dim(H^{(l)})$ is an orthonormal basis in $H^{(l)}$ and U is a unitary operator acting on $H^{(l)}$. A direct calculation of $tr(\rho\Pi^{(l)}\rho)$ and comparison with $\rho^{(l)} = tr_{\bar{l}}(\rho)$ gives, assuming that $\{U|k_l\rangle\}$ is the eigenbasis of $\rho^{(l)}$, $tr(\rho\Pi^{(l)}(\rho)) = \sum_{k_l}(\langle k_l|U^\dagger\rho^{(l)}U|k_l\rangle)^2 = \sum_{k_l}\lambda_{k_l}^2 = tr(\rho^{(l)})^2$, where $\{\lambda_{k_l}\}$ are the eigenvalues of $\rho^{(l)}$.

From the definition of $N_l(\rho)$ (Eq.(1)) we get

$$N_l(\rho) = \frac{d_l}{d_l - 1}||\rho||^2 - \min_{\Pi^{(l)}}(2tr(\rho\Pi^{(l)}(\rho)) - ||\Pi^{(l)}(\rho)||^2).$$

For a pure state $||\rho||^2 = 1$ and $||\Pi^{(l)}(\rho)||^2 = tr(\rho\Pi^{(l)}(\rho))$ so that

$$N_l(\rho) = \frac{d_l}{d_l - 1}(1 - \min_{\Pi^{(l)}} tr(\rho\Pi^{(l)}(\rho))).$$

The minimum is over the von-Neumann measurements leaving the marginal state $\rho^{(l)}$ invariant, that is $\sum_k \Pi_k^{(l)}\rho^{(l)}\Pi_k^{(l)} = \rho^{(l)}$, or,

$$\sum_{k_l}\langle k_l|U^\dagger\rho^{(l)}U|k_l\rangle U|k_l\rangle\langle k_l|U^\dagger = \rho^{(l)}.$$

This is the spectral decomposition of $\rho^{(l)}$ which is consistent with our choice of $\{U|k_l\rangle$ to be the eigenbasis of $\rho^{(l)}$. Since $tr(\rho\Pi^{(l)}(\rho))$ is simply the trace of $(\rho^{(l)})^2$, the minimization in the definition of N_l Eq.(1) drops out and we get

$$N_l(|\psi\rangle\langle\psi|) = \frac{d_l}{d_l - 1}(1 - tr(\rho^{(l)})^2).$$

Corollary: For an n-partite pure state $\rho = |\psi\rangle\langle\psi|$

$$D_l(\rho) = N_l(\rho) \tag{3}$$

where $D_l(\rho)$ is the geometric discord of ρ with von-Neomann measurement on the lth part [6]. This important result follows trivially, because $D_l(\rho)$ requires maximization over all von-Neumann measurements on the lth part which is obtained if the $\{U|k_l\rangle\}$ forms the eignbasis of $\rho^{(l)}$.

It is interesting to compare $N_l(\rho)$ (Eq.(1)) with measures of entanglement of pure multipartite states. For a bipartite pure state ρ_{AB} we have, for the concurrence,

$$C(\rho_{AB}) = \sqrt{2(1 - tr(\rho_A^2)}$$

which is related to $N_l(\rho_{AB})$ by

$$N_l(\rho_{AB}) = \frac{d_l}{2(d_l-1)}C^2(\rho_{AB}).$$

Thus, for pure bipartite states, non-locality is simply related to concurrence.

The Meyer-Wallach measure of entanglement of multipartite pure states is

$$Q(|\psi\rangle) = \frac{1}{n}\sum_{k=1}^{n} 2(1 - tr(\rho_k^2))$$

where ρ_k is the reduced density operator for the kth part. Thus,

$$Q(|\psi\rangle) = \frac{2}{n}\sum_{l=1}^{n}\left(\frac{d_l-1}{d_l}\right) N_l(|\psi\rangle\langle\psi|).$$

Thus the Meyer-Wallach measure of pure state multipartite entanglement is the average of non-locality over the parts of the system.

Non-locality in the multipartite mixed states: To get $N_l(\rho)$ in this case, we start with the Bloch representation of a multipartite state ρ[8]. Bloch representation [8] of a n-partite density operetor is

$$\rho = \frac{1}{\Pi_k^n d_k}\{\otimes_k^n I_{d_k} + \sum_{k\in\mathcal{N}}\sum_{\alpha_k} s_{\alpha_k}\lambda_{\alpha_k}^{(k)} + \sum_{2\le M\le n}\ \sum_{\{k_1,k_2,\cdots,k_M\}}\ \sum_{\alpha_{k_1}\alpha_{k_2}\cdots\alpha_{k_M}} \tilde{t}_{\alpha_{k_1}\alpha_{k_2}\cdots\alpha_{k_M}}\lambda_{\alpha_{k_1}}^{(k_1)}\lambda_{\alpha_{k_2}}^{(k_2)}\cdots\lambda_{\alpha_{k_M}}^{(k_M)}\}. \quad (4)$$

where $\mathcal{N} = \{1, 2, \cdots, n\}$ and

$$\lambda_{\alpha_{k_1}}^{(k_1)} = (I_{d_1}\otimes I_{d_2}\otimes\cdots\otimes\lambda_{\alpha_{k_1}}\otimes I_{d_{k_1+1}}\otimes\cdots\otimes I_{d_n})$$

$$\lambda_{\alpha_{k_2}}^{(k_2)} = (I_{d_1}\otimes I_{d_2}\otimes\cdots\otimes\lambda_{\alpha_{k_2}}\otimes I_{d_{k_2+1}}\otimes\cdots\otimes I_{d_n})$$

$$\lambda_{\alpha_{k_1}}^{(k_1)}\lambda_{\alpha_{k_2}}^{(k_2)} = (I_{d_1}\otimes I_{d_2}\otimes\cdots\otimes\lambda_{\alpha_{k_1}}\otimes I_{d_{k_1+1}}\otimes\cdots\otimes\lambda_{\alpha_{k_2}}\otimes I_{d_{k_2+1}}\otimes I_{d_n}) \quad (5)$$

$\mathbf{s}^{(k)}$ is a Bloch vector corresponding to kth subsystem, $\mathbf{s}^{(k)} = [s_{\alpha_k}]_{\alpha_k=1}^{d_k^2-1}$ and

$$\tilde{t}_{\alpha_{k_1}\alpha_{k_2}\ldots\alpha_{k_M}} = \frac{d_{k_1}d_{k_2}\ldots d_{k_M}}{2^M} tr[\rho\lambda_{\alpha_{k_1}}^{(k_1)}\lambda_{\alpha_{k_2}}^{(k_2)}\cdots\lambda_{\alpha_{k_M}}^{(k_M)}]. \quad (6)$$

For more details see ref.[8–10].

Recently, for a bipartite system ab ($n = 2$) with states in $\mathcal{H}^a\otimes\mathcal{H}^b$, $dim(\mathcal{H}^a) = d_a$, $dim(\mathcal{H}^b) = d_b$, S. Luo and S. Fu introduced the following generic expression for Min [2]

$$N_a(\rho) = tr(TT^t) - \min_A tr(ATT^tA^t), \quad (7)$$

where $T = [t_{ij}]$ is an $d_a^2\times d_b^2$ matrix and the minimum is taken over all $d_a\times d_a^2-1$-dimensional isometric matrices $A = [a_{ji}]$ such that $a_{ji} = tr(|j\rangle\langle j|X_i) = \langle j|X_i|j\rangle$, $j = 1, 2, \ldots, d_a$; $i = 1, 2, \ldots, d_a^2 - 1$ and $\{|j\rangle\}$ is any orthonormal basis in $\mathcal{H}^a$. we generalize this result to n-partite quantum states, in theorem 2 and 3.

Theorem 2. Let $\rho_{12\cdots n}$ be a n-partite state defined by Eq.(4), then

$$N_l(\rho)=\frac{d_l}{(d_l-1)\Pi_k^n d_k}\{\sum_{1\le M\le n-1}\sum_{\{k_1,k_2,\cdots,k_M\}\subseteq\mathcal{N}-\{l\}}\frac{d_l d_{k_1}d_{k_2}\cdots d_{k_M}}{2^{M+1}}||\mathcal{T}^{\{l,k_1,k_2,\cdots,k_M\}}||^2-\min_{A^{(l)}} tr(A^{(l)}K^{(l)}(A^{(l)})^t)\}, \tag{8}$$

where the $(d_l^2-1)\times(d_l^2-1)$ symmetric matrix $K^{(l)}$ is defined as

$$K^{(l)}_{\alpha_l\beta_l}=\sum_{1\le M\le n-1}\sum_{\{k_1,k_2,\cdots,k_M\}\subseteq\mathcal{N}-\{l\}}\sum_{\alpha_{k_1}\alpha_{k_2}\cdots\alpha_{k_M}}\frac{d_l d_{k_1}d_{k_2}\cdots d_{k_M}}{2^{M+1}}t_{\alpha_l\alpha_{k_1}\alpha_{k_2}\cdots\alpha_{k_M}}t_{\beta_l\alpha_{k_1}\alpha_{k_2}\cdots\alpha_{k_M}}.$$

$\mathcal{T}=[t_{i_1i_2\cdots i_M}]=[tr(\rho\lambda^{(k_1)}_{\alpha_{k_1}}\lambda^{(k_2)}_{\alpha_{k_2}}\cdots\lambda^{(k_M)}_{\alpha_{k_M}})]$, and the maximum is taken over all $d_l\times(d_l^2-1)$ dimensional matrices $A^{(k)}=[a_{ji_l}]$, such that $a_{ji_l}=tr(|j\rangle\langle j|\frac{\lambda^{(l)}_{i_l}}{\sqrt{2}}),\ j=1,2,\ldots,d_l;\ i_l=1,2,\ldots,d_l^2-1$ and $\{|j\rangle\}$ is any orthonormal basis for $\mathcal{H}^{(l)}$. In particular, we have

$$N_l(\rho)\le\sum_{i=1}^{d_l^2-d_l}\eta_i, \tag{9}$$

where$\{\eta_i\ :\ i=1,2,\cdots,d_l^2-1\}$ are the eigenvalues of the $(d_l^2-1)\times(d_l^2-1)$ symmetric matrix $K^{(l)}$ listed in non-increasing order. Furthermore, if $\rho^{(l)}=tr_{\bar{l}}\rho_{12\cdots n}$ is non-degenerate with spectral projections $\{|j\rangle\langle j|\}$, then

$$N_l(\rho)=\frac{d_l}{(d_l-1)\Pi_k^n d_k}\{\sum_{1\le M\le n-1}\sum_{\{k_1,k_2,\cdots,k_M\}\subseteq\mathcal{N}-\{l\}}\frac{d_l d_{k_1}d_{k_2}\cdots d_{k_M}}{2^{M+1}}||\mathcal{T}^{\{l,k_1,k_2,\cdots,k_M\}}||^2-tr(A^{(l)}K^{(l)}(A^{(l)})^t)\}, \tag{10}$$

Theorem 3. If the lth part of a n-partite quantum system is a qubit ($d_l=2$), then

$$N_l(\rho)=\frac{d_l}{(d_l-1)\Pi_k^n d_k}\left[\sum_{1\le M\le n-1}\sum_{\{k_1,\ldots,k_M\}\subseteq\mathcal{N}-\{l\}}\frac{d_{k_1}d_{k_2}\cdots d_{k_M}}{2^M}||\mathcal{T}^{\{l,k_1,\ldots,k_M\}}||^2-\begin{cases}\frac{\mathbf{s}^{(l)t}K^{(l)}\mathbf{s}^{(l)}}{||\mathbf{s}^{(l)}||^2}, & \text{if } \mathbf{s}^{(l)}\neq 0\\ \eta_{min}, & \text{if } \mathbf{s}^{(l)}=0\end{cases}\right] \tag{11a}$$

where $\mathbf{s}^{(l)}$ is the coherent vector of $\rho^{(l)}$ and η_{min} is the smallest eigenvalue of the matrix $K^{(l)}$ which is a 3×3 real symmetric matrix, defined as

$$K^{(l)}=\sum_{1\le M\le n-1}\sum_{\{k_1,\ldots,k_M\}\subseteq\mathcal{N}-\{l\}}\sum_{\alpha_{k_1}\alpha_{k_2}\cdots\alpha_{k_M}}\frac{d_{k_1}d_{k_2}\cdots d_{k_M}}{2^M}t_{\alpha_l\alpha_{k_1}\alpha_{k_2}\cdots\alpha_{k_M}}t_{\beta_l\alpha_{k_1}\alpha_{k_2}\cdots\alpha_{k_M}}, \tag{11}$$

For n-qubit $(d_i=2,\ i=1,2,\cdots,n)$,

$$N_l(\rho)=\frac{1}{2^{(n-1)}}\left[\sum_{1\le M\le n-1}\sum_{\{k_1,\ldots,k_M\}\subseteq\mathcal{N}-\{l\}}||\mathcal{T}^{\{l,k_1,\ldots,k_M\}}||^2-\begin{cases}\frac{\mathbf{s}^{(l)t}K^{(l)}\mathbf{s}^{(l)}}{||\mathbf{s}^{(l)}||^2}, & \text{if } \mathbf{s}^{(l)}\neq 0\\ \eta_{min}, & \text{if } \mathbf{s}^{(l)}=0\end{cases}\right]$$

and

$$K^{(l)}=\sum_{1\le M\le n-1}\sum_{\{k_1,\ldots,k_M\}\subseteq\mathcal{N}-\{l\}}\sum_{\alpha_{k_1}\alpha_{k_2}\cdots\alpha_{k_M}}t_{\alpha_l\alpha_{k_1}\alpha_{k_2}\cdots\alpha_{k_M}}t_{\beta_l\alpha_{k_1}\alpha_{k_2}\cdots\alpha_{k_M}}, \tag{12}$$

The proofs of theorems 2 and 3 is a straightforward generalization of those of theorems 2 and 3 respectively in ref. [2] to the multipartite case, so that we skip these proofs.

Relation between the non-locality and geometric quantum discord for arbitrary n-qubit states : We saw (see(Eq.(2))) that the non-locality and geometric discord are equal for arbitrary n-partite pure states. In this section we find a class of general n-qubit states for which these quantities coincide. Consider a n-qubit state ρ. The geometric discord for such a state corresponding to the von-Neumann measurement on lth qubit is given by

$$D_l(\rho) = \frac{1}{2^{(n-1)}}\left[||s^{(l)}||^2 + \sum_{1\leq M\leq n-1}\sum_{\{k_1,\dots,k_M\}\subseteq\mathcal{N}-\{l\}} ||\mathcal{T}^{\{l,k_1,\dots,k_M\}}||^2 - \lambda_{max},\right] \tag{13}$$

where $s^{(l)}$ is the coherent vector of $\rho^{(l)}$ (reduced density operator for the lth part), $\mathcal{T} = [t_{\alpha_{k_1}\alpha_{k_2}\dots\alpha_{k_M}}] = [tr(\rho\lambda^{(k_1)}_{\alpha_{k_1}}\lambda^{(k_2)}_{\alpha_{k_2}}\cdots\lambda^{(k_M)}_{\alpha_{k_M}})]$, and λ_{max} is the largest eigenvalue of the 3×3 real symmetric matrix

$$G^{(l)} = \mathbf{s}^{(l)}(\mathbf{s}^{(l)})^t + K^{(l)} \tag{14}$$

where $K^{(l)}$ is given by Eq.(12) for n qubits. The non-locality for the n-qubit state ρ is given by Eq.(12). We now consider two cases

Case I : $\mathbf{s}^{(l)} \neq 0$. By Eq.(14) we get

$$\hat{e}^t G^{(l)}\hat{e} = \hat{e}^t\mathbf{s}^{(l)}\mathbf{s}^{(l)}\hat{e} + \hat{e}^t K^{(l)}\hat{e}$$

where $\hat{e}\in R^3$ is an arbitrary unit vector, choosing $\hat{e} = \frac{\mathbf{s}^{(l)}}{||\mathbf{s}^{(l)}||}$, we get

$$\frac{(\mathbf{s}^{(l)})^t K^{(l)}\mathbf{s}^{(l)}}{||\mathbf{s}^{(l)}||^2} = \frac{(\mathbf{s}^{(l)})^t G^{(l)}\mathbf{s}^{(l)}}{||\mathbf{s}^{(l)}||^2} - ||\mathbf{s}^{(l)}||^2$$

Substituting in Eq.(12) we get

$$N_l(\rho) = \frac{1}{2^{(n-1)}}\left[||\mathbf{s}^{(l)}||^2 + \sum_{1\leq M\leq n-1}\sum_{\{k_1,\dots,k_M\}\subseteq\mathcal{N}-\{l\}} ||\mathcal{T}^{\{l,k_1,\dots,k_M\}}||^2 - \frac{{\mathbf{s}^{(l)}}^t G^{(l)}\mathbf{s}^{(l)}}{||\mathbf{s}^{(l)}||^2}\right] \tag{15}$$

If $\frac{\mathbf{s}^{(l)}}{||\mathbf{s}^{(l)}||}$ is the eigenvector of $G^{(l)}$ with the largest eigenvalue then the right hand side of Eq.(14) gives the geometric discord $D_l(\rho)$ so that under this condition $N_l(\rho) = D_l(\rho)$. The above condition can be equivalently stated as

$$[\mathbf{s}^{(l)}(\mathbf{s}^{(l)})^t, K(l)] = 0$$

and

$$||\mathbf{s}^{(l)}||^2 + \eta_l \geq \eta_{i\neq l}$$

where $\{\eta_i\}$ are the eigenvalues of $K^{(l)}$ and η_l is the eigenvalue corresponding to the eigenvector $\frac{\mathbf{s}^{(l)}}{||\mathbf{s}^{(l)}||}$.

Case II: $\mathbf{s}^{(l)} = 0$. In this case ρ has one doubly degenerate eigenvalue. With $\mathbf{s}^{(l)} = 0$ we get from Eq.(14)

$$\hat{e}^t G^{(l)}\hat{e} = \hat{e}^t K^{(l)}\hat{e}. \tag{16}$$

To get non-locality we have to minimize the right hand side while the geometric discord requires maximization of the left hand side. Under these conditions, the equality in Eq.(15) is preserved if $G^{(l)} = K^{(l)}$ has a single three-fold degenerate eigenvalue,($\eta_1 = \eta_2 = \eta_3$). Thus when $\mathbf{s}^{(l)} = 0$,



Figure 1 : The variation of Min $N_l(\rho(p))$ (dashed carve) and the geometric discord $D_l(\rho(p))$ (continuous carve) for the state as in (a) Eq(17) (b) Eq(18) (c) Eq(19) (d) Eq(20) with parameter p.

$N_l(\rho) = D_l(\rho)$ provided the matrix $K^{(l)}$ has a single three fold degenerate eigenvalue.

Examples : As our first example we consider the set of three qubit states comprising the convex combination of $|GHZ\rangle = \frac{1}{\sqrt{2}}(|000\rangle + |111\rangle)$ and $|W\rangle = \frac{1}{\sqrt{3}}(|001\rangle + |010\rangle + |100\rangle)$,

$$\rho(p) = p|GHZ\rangle\langle GHZ| + (1-p)|W\rangle\langle W|. \tag{17}$$

The $K^{(1)}$ matrix of this state is

$$K^{(1)} = \mathrm{diag}[2p^2 + \frac{16}{9}(1-p)^2, 2p^2 + \frac{16}{9}(1-p)^2, 2p^2 + \frac{19}{9}(1-p)^2 - \frac{4}{3}p(1-p)]$$

with the coherent vector for the first qubit

$$\mathbf{s}^{(1)} = [0, 0, \frac{1}{3}(1-p)]^t \neq 0$$

so that case I applies. We find that $[\mathbf{s}^{(1)}(\mathbf{s}^{(1)})^t, K^{(1)}] = 0$ and the condition $||\mathbf{s}^{(1)}||^2 + \eta_1 \geq \eta_{i\neq 1}$, ($\eta_1$ is the eigenvalue of $K^{(1)}$ matrix corresponding to eigenvector $\frac{\mathbf{s}^{(1)}}{||\mathbf{s}^{(1)}||}$), is satisfied when $p \leq \frac{1}{4}$ and $p = 1$. This is depicted in fig.(1a).

The second example consists of

$$\rho(p) = p|\widetilde{W}\rangle\langle\widetilde{W}| + (1-p)|W\rangle\langle W| \tag{18}$$

where $|\widetilde{W}\rangle$ is the flipped $|W\rangle$ state, $\sigma_x \otimes \sigma_x \otimes \sigma_x|W\rangle$. The $K^{(1)}$ matrix of this state is

$$K^{(1)} = \mathrm{diag}[\frac{16}{9}p^2 + \frac{16}{9}(1-p)^2, \frac{16}{9}p^2 + \frac{16}{9}(1-p)^2, \frac{19}{9}p^2 + \frac{19}{9}(1-p)^2 - \frac{10}{3}p(1-p)]$$

with the coherent vector for the first qubit

$$\mathbf{s}^{(1)} = [0, 0, \frac{1}{3}(1-2p)]^t \neq 0,$$

so that case I applies. We find that $[\mathbf{s}^{(1)}(\mathbf{s}^{(1)})^t, K^{(1)}] = 0$ and the condition $||\mathbf{s}^{(1)}||^2 + \eta_1 \geq \eta_{i\neq 1}$, is satisfied when $p \leq 0.1127$ and $p \geq 0.8873$. The results are shown in fig.(1b).

The third example consists of

$$\rho(p) = p|GHZ\rangle\langle GHZ| + (1-p)|GHZ_-\rangle\langle GHZ_-|. \tag{19}$$

where $|GHZ_-\rangle = \frac{1}{\sqrt{2}}(|000\rangle - |111\rangle)$. The $K^{(1)}$ matrix of this state is

$$K^{(1)} = \mathrm{diag}[2(2p^2-1)^2, 2(2p^2-1)^2, 2(2p^2-1)^2]$$

and the coherent vector for the first qubit

$$\mathbf{s}^{(1)} = 0,$$

so that case II applies. $K^{(1)}$ does not have a single triply degenerate eigenvalue, for all p, except $p = 0$ and $p = 1$. Therefore $N_l(\rho) \neq D_l(\rho)$ for all p between 0 and 1. The results are shown in fig.(1c).

The last example consists of the states

$$\rho(p) = p|GHZ\rangle\langle GHZ| + (1-p)|GHZ_1\rangle\langle GHZ_1|. \tag{20}$$

where $|GHZ_1\rangle = \frac{1}{\sqrt{2}}(|001\rangle + |110\rangle)$. The $K^{(1)}$ matrix of this state is

$$K^{(1)} = \mathrm{diag}[2(p^2-(1-p)^2), 2(p^2-(1-p)^2), 2(p^2-(1-p)^2)]$$

and the coherent vector for the first qubit

$$\mathbf{s}^{(1)} = 0,$$

so that case II applies. $K^{(1)}$ does have a single triply degenerate eigenvalue, for all p. Therefore $N_l(\rho) = D_l(\rho)$ for all p as shown in fig.(1d).

The conditions obtained in case I and case II above define a class of states for which MiN and the geometric quantum discord coincide. Recently quantum discord is shown to measure the quantumness of the state rather than genuine quantum correlation [13]. Thus for the class of states with equal $N_l(\rho)$ and $D_l(\rho)$, MiN seems to be identical or simply related to quantumness of the state. An understanding of relation between quantumness and non-locality by other routes will then be interesting. At any rate, it is interesting to explore the relation between $N_l(\rho)$ and $D_l(\rho)$ in the states for which they do not coincide, because these will improve our understanding of quantum correlations.

Acknowledgments : This work was supported by the BCUD grant RG-13. ASMH acknowledges University of Pune for hospitality during his visit when this work was carried out.

---

[1] L. B. Fu, Europhys. Lett. **75**, 1 (2006).
[2] S. Luo and S. Fu, Phys. Rev. Lett. **106**, 120401 (2012).
[3] H. Ollivier and W. H. Zurek, Phys. Rev. Lett. **88**, 017901 (2001).
[4] L. Henderson and V. Vedral, J. Phy. A **34**, 6899 (2001).

[5] B. Dakic, V. Vedral, and C. Brukner, Phys. Rev. Lett. **105**,190502 (2010).
[6] Ali Saif M. Hassan and Pramod S. Joag, J. Phys. A: Math. Theor. **45**, 345301 (2012).
[7] D. Girolami and G. Adesso, Phys. Rev. A 84, 052110 (2011).
[8] Ali Saif M. Hassan and Pramod S. Joag, Quantum Inf. Comput. **8**, 773 (2008).
[9] G. Mahler, Volker A. Weberruss, *Quantum networks* (Springer-Verlag Berlin Heidelberg 1995). **8**, 773 (2008).
[10] M. S. Byrd and n. Khaneja, Phys. Rev. A **68**, 062322 (2003); G. Kimura, Phys. Lett. A **314**, 339 (2003).
[11] T. G. Kolda, "Multilinear operator for higher order decompositions", Tech. Report SAnD2006-2081, Sandia national Laboratories, Albuquerque, new Mexico and Livermore, California Apr. 2006.
[12] L. De Lathauwer, B. De Moor and J. Vandewalle, *SIAM J. Matrix Anal. A.* **21**, 1253 (2000).
[13] M. Gessner, Elsi-Mari Laine, Heinz-Peter Breuer, J. Piilo, Phys. Rev. A **85**, 052122 (2012).